

Department of Mathematics and Statistics


Preprint MPS-2016-06

15 April 2016

$\mathcal{D}$

∞

Nikos Katzourakis



D-SOLUTIONS TO THE SYSTEM OF VECTORIAL CALCULUS OF VARIATIONS IN L^1 VIA THE BAIRE CATEGORY METHOD FOR THE SINGULAR VALUES

GISELLA CROCE, NIKOS KATZOURAKIS AND GIOVANNI PISANTE

Abstract. For $H \in C^2(\mathbb{R}^{N \times n})$ and $u : \mathbb{R}^n \rightarrow \mathbb{R}^N$, consider the system

$$(1) \quad H_P - \mathbb{H}_P + H[H_P]^\perp H_{PP} (Du) : D^2 u = 0 :$$

The PDE system (1) is associated to the supremal functional

$$(2) \quad E_1(u; \emptyset) = \int_{\emptyset} H(Du) dx ; \quad u \in W_{loc}^{1,1}(\emptyset; \mathbb{R}^N); \quad \emptyset \subset \mathbb{R}^n ;$$

and first arose in recent work of the 2nd author as the analogue of the Euler-Lagrange equation. Herein we employ the Dacorogna-Marcellini Baire Category method to construct D-solutions to the Dirichlet problem for (1), an apt notion of generalised solutions recently proposed for fully nonlinear systems. Our D-solutions to (1) are $W^{1,1}$ -submersions corresponding to "critical points" of (2) and are obtained without any convexity hypotheses. Along the way we establish a result of independent interest by proving existence of strong solutions to the singular value problem for general dimensions $n \in \mathbb{N}$.

1. Introduction

Let $H \in C^2(\mathbb{R}^{N \times n})$ be a given function and $\Omega \subset \mathbb{R}^n$ a given open set, $n, N \geq 2$. In this paper we are interested in the problem of existence of appropriately defined generalised solutions with given Dirichlet boundary conditions to the second order PDE system

$$(1.1) \quad A_1 u := H_P - H_P + H[H_P]^\perp H_{PP} (Du) : D^2 u = 0 :$$

In the above, the subscript P denotes the derivative of H with respect to its matrix variable, while

$$\begin{aligned} Du(x) &= (D_i u(x))_{i=1, \dots, n}^{j=1, \dots, N} \in \mathbb{R}^{N \times n}; \\ D^2 u(x) &= (D_{ij}^2 u(x))_{i,j=1, \dots, n}^{j=1, \dots, N} \in \mathbb{R}_s^{N \times n^2}; \end{aligned}$$

denote respectively the gradient matrix and the hessian tensor of (smooth) maps $u : \mathbb{R}^n \rightarrow \mathbb{R}^N$. The notation $[H_P]^\perp$ symbolises the orthogonal projection on the orthogonal complement of the range of the linear map $\mathbb{H}(P) : \mathbb{R}^n \rightarrow \mathbb{R}^N$:

$$(1.2) \quad [H_P(P)]^\perp := \text{Proj}_{R(H_P(P))^\perp} :$$

Key words and phrases. Vectorial Calculus of Variations in L^1 ; Generalised solutions; Fully nonlinear systems; 1-Laplacian; Young measures; Singular Value Problem, Baire Category method; Convex Integration.

In index form, (1.1) reads

$$\sum_{i,j=1}^N \sum_{k,l=1}^N H_{P_{ij}}(Du) H_{P_{kl}}(Du) + H(Du) \sum_{i,j=1}^N H_{P_{ij}}(Du) - D_{ij}^2 u = 0;$$

where $i, j, k, l = 1, \dots, N$. Our general notation is either self-explanatory or a convex combination of standard symbolisations as e.g. in [E, D, EG, DM2]. The system (1.1) is the analogue of the Euler-Lagrange equation when one considers nonstandard vectorial variational problems in the space L^1 for the supremal functional

$$(1.3) \quad E_1(u; \varphi) := \int_{\Omega} H(Du) \, dx; \quad u \in W_{loc}^{1,1}(\Omega; \mathbb{R}^N); \quad \varphi \in \mathcal{B}$$

and first arose in recent work of the second author ([K1]). Calculus of Variations in L^1 , as the field is known today, was initiated by G. Aronsson in the 1960s who studied the scalar case $N = 1$ quite systematically ([A1]–[A7]). Since then the area has been developed marvellously due to both the intrinsic mathematical interest and the importance for applications. In particular, the theory of Viscosity Solutions of Crandall-Ishii-Lions played a fundamental role in the study of the generally singular solutions to the scalar version of (1.1). When $N = 1$, the respective single equation simplifies to

$$(1.4) \quad A_1 u = H_P(Du) - H_P(Du) : D^2 u = 0$$

and is known as the "Aronsson equation". For a pedagogical introduction to the scalar case with numerous references see e.g. [K7, C]. See also [Pi] for a comparison between the Viscosity Solutions approach and the Baire Category one.

On

case of $\text{rank}(Du) = 2$, the appropriate minimality notion connecting (1.1) to (1.3) is not the obvious extension of Aronsson's scalar notion of Absolute Minimiser (see [K2, K10, K13]).

Perhaps the greatest difficulty associated to the study of (1.1) is that it is quasi-linear, non-divergence and non-monotone and all standard approaches in order to define generalised solutions based on integration-by-parts or on the maximum principle seem to fail. Motivated partly by the systems arising in Calculus of Variations in L^1 , the second author has recently proposed in [K8] a new efficient theory of

of Du in the Young measures valued into the sphere $\overline{R}_s^{N-n^2}$:

$$D^{1,h}_m Du \ast D^2u; \quad \text{in } Y; \quad \overline{R}_s^{N-n^2}; \quad \text{as } m \rightarrow 1: \quad$$

Our main ingredient for the solvability of (1.11) is a result of independent interest about the solvability of the following fully non-linear system, usually referred to as the prescribed singular value problem

$$(1.13) \quad \begin{aligned} \lambda_i(Du) &= 1; \quad \text{a.e. on } \Omega; \quad i = 1, \dots, n \wedge N; \\ u &= g; \quad \text{on } \partial\Omega. \end{aligned}$$

In (1.13), $\lambda_1(Du), \dots, \lambda_{n \wedge N}(Du)$ denotes the set of singular values of the matrix Du , namely the eigenvalues of the matrix $(Du^T Du)^{1/2}$ in increasing order and we symbolise $n \wedge N = \min\{n, N\}$.

Systems of PDEs involving singular values, mostly related to non-convex problems in Calculus of Variations, have been considered by several authors (cf. for instance [Cr, BCR, DR, DPR]). In particular the problem of finding sufficient conditions on the boundary datum g in order to get existence of solutions to problem (1.13) has been addressed, in the special case $n = N$, in [DM1, DM2, DT]. To the best of our knowledge no results are known for the case $n \neq N$. Accordingly, we establish the following result.

Theorem 5. Let $\Omega \subset \mathbb{R}^n$ be an open set. Assume that $g \in W^{1,1}(\Omega; \mathbb{R}^N)$ is such that $\lambda_{n \wedge N}(Dg) < 1$, a.e. on Ω . Then, there exists an infinite set of solutions $u \in W^{1,1}_g(\Omega; \mathbb{R}^N)$ to the system (1.13).

The proof of the previous theorem can be obtained as an application of the general existence theory for differential inclusion via the Baire Category method (cf. [PD]), in the same spirit as in the $N = n$ case. It relies on the characterisation of the rank-one convex envelope of the set of matrices

$$E := \left\{ Q \in \mathbb{R}^{N \times n} : \lambda_i(Q) = 1; \quad i = 1, \dots, n \wedge N \right\};$$

which is

$$\text{Rco}E = \left\{ Q \in \mathbb{R}^{N \times n} : \lambda_{n \wedge N}(Q) \leq 1 \right\};$$

as proved in Theorem 11.

In order to address the question of the existence of solutions to the problem (1.11) with $g \in W^{1,1}(\Omega; \mathbb{R}^N)$, some comments on the admissible regularity of the boundary datum in Theorem 5 are in order. Indeed, the piecewise a.e. regularity of the datum g can be weakened to Lipschitz continuity if we restrict slightly the bound on the norm $\lambda_{n \wedge N}(Dg)$. This result, precisely stated in Corollary 6 that follows, is a simple consequence of the convexity of the rank-one convex hull $\overline{\text{co}}^{\text{r.o.}}$ (cf. Theorem 11) and of the approximation result proved in [DM2, Corollary 10.21].

Corollary 6. Let $\Omega \subset \mathbb{R}^n$ be an open set. Assume that $g \in W^{1,1}(\Omega; \mathbb{R}^N)$ is such that for some $\varepsilon > 0$ we have $\lambda_{n \wedge N}(Dg) \leq 1 - \varepsilon$, a.e. on Ω . Then there exists a infinite set of solutions $u \in W^{1,1}_g(\Omega; \mathbb{R}^N)$ to the system (1.13).

The rest of the paper is organised as follows. In the next section we recall some known results about Young measures valued into spheres. In Section 3 we provide the proof of the existence of solutions for the prescribed singular value problem and in the last section we prove existence and geometric properties of solutions to the problem (1.10).

2. Young measures valued into spheres

Here we collect some basic material taken from [K8] which can be found in different guises and in greater generality e.g. in [CFV, FG]. Let $E \subset \mathbb{R}^n$ be measurable and consider the L^1 space of strongly measurable maps valued in the continuous functions over the sphere $\overline{\mathbb{R}}_s^{N-n^2}$ (for details on these spaces see e.g. [Ed, FL]):

$$L^1(E; C(\overline{\mathbb{R}}_s^{N-n^2})) :$$

This space consists of Carathéodory functions $\gamma : E \rightarrow \overline{\mathbb{R}}_s^{N-n^2}$ for which

$$\|\gamma\|_{L^1(E; C(\overline{\mathbb{R}}_s^{N-n^2}))} = \int_E \left(\int_{\overline{\mathbb{R}}_s^{N-n^2}} \langle \gamma(x), \xi \rangle d\xi \right) dx < \infty :$$

The dual of this (separable) Banach space is

$$L^1_w(E; M(\overline{\mathbb{R}}_s^{N-n^2})) = L^1(E; M(\overline{\mathbb{R}}_s^{N-n^2})) :$$

The dual space above consists of measure-valued maps $\mu : E \rightarrow \mathcal{M}(\overline{\mathbb{R}}_s^{N-n^2})$ which are weakly* measurable, that is, for any fixed $\phi \in C(\overline{\mathbb{R}}_s^{N-n^2})$, the function

$$E \ni x \mapsto \int_{\overline{\mathbb{R}}_s^{N-n^2}} \phi(\xi) d\mu_x(\xi) \text{ is measurable,} \quad \text{[Ed, FL]}$$

3. The prescribed singular value problem

In this section we prove Theorem 5 and Corollary 6. To this aim we start by recalling for the convenience of the reader some well known results about generalised convex hulls of sets of matrices (for further details we refer to the books [DM2] and [D]).

For $Q \in \mathbb{R}^{N \times n}$ we set

$$T(Q) := \begin{pmatrix} Q; \text{adj}_2 Q; \dots; \text{adj}_{N \wedge n} Q \end{pmatrix} \in \mathbb{R}^{(N;n)};$$

where $\text{adj}_s Q$ stands for the matrix of all $s \times s$ subdeterminants of the matrix Q , $1 \leq s \leq N \wedge n = \min\{N, n\}$ and

$$(N;n) := \sum_{s=1}^{N \wedge n} \binom{N}{s} \binom{n}{s}; \quad \binom{N}{s} = \frac{N!}{s!(N-s)!}.$$

Definition 7. Consider a function $f : \mathbb{R}^{N \times n} \rightarrow \mathbb{R} \cup \{+\infty\}$.

- (1) f is said to be polyconvex if there exists a convex function $g : \mathbb{R}^{(N;n)} \rightarrow \mathbb{R} \cup \{+\infty\}$ such that $f(Q) = g(T(Q))$.
- (2) f is said to be rank one convex if

$$f(Q + (1-t)R) \leq t f(Q) + (1-t) f(R)$$

for every $t \in [0, 1]$ and every $Q, R \in \mathbb{R}^{N \times n}$ with $\text{rk}(R - Q) = 1$.

It is well known that if a function is polyconvex, then it is rank one convex. Next we recall the corresponding notions of convexity for sets.

Definition 8. Let E be a subset of $\mathbb{R}^{N \times n}$.

- (1) We say that E is polyconvex if there exists a convex set $S \subset \mathbb{R}^{(N;n)}$ such that $E = T^{-1}(S)$.

Analogous representations to (32

such that $(Z)_{n \wedge N; n \wedge N} = 1$ and the other entries of Z are null. Then $Q + \epsilon Z$ would

4. D-solutions to the PDE system arising in vectorial Calculus of Variations in L^1

In this section we establish the proofs of Theorem 3 and of Proposition 4 by utilising Corollary 6

Now we complete the proof of the proposition. By our assumptions on H (see Theorem 3), for any u as above we have

$$H(Du) = h(Du)Du^\top = h(c^2)I$$

and by splitting the identity as

$$c^2 I = [c]_j [O]_j [c]_j [O]_j^\top$$

where $[c]_j [O]_j \in \mathbb{R}^{N \times N} \subset \mathbb{R}^N$

for a.e. $x \in \mathbb{R}^2$. Since $D u \in L^1(\mathbb{R}^2)$

References

- AK. H. Abugirda, N. Katzourakis, Existence of 1D vectorial Absolute Minimisers in L^1 under minimal assumptions , ArXiv preprint, <http://arxiv.org/pdf/1604.03068.pdf> .
- AM. L. Ambrosio, J. Maj, Very weak notions of differentiability , Proceedings of the Royal Society of Edinburgh A 137 (2007), 447 - 455.
- A1. G. Aronsson, Minimization problems for the functional $\sup_x F(x; f(x); f^0(x))$, Arkiv för Mat. 6 (1965), 33 - 53.
- A2. G. Aronsson, Minimization problems for the functional $\sup_x F(x; f(x); f^0(x))$ II , Arkiv för Mat. 6 (1966), 409 - 431.
- A3. G. Aronsson, Extension of functions satisfying Lipschitz conditions , Arkiv för Mat. 6 (1967), 551 - 561.
- A4. G. Aronsson, On the partial differential equation $u_x^2 u_{xx} + 2 u_x u_y u_{xy} + u_y^2 u_{yy} = 0$, Arkiv för Mat. 7 (1968), 395 - 425.
- A5. G. Aronsson, Minimization problems for the functional $\sup_x F(x; f(x); f^0(x))$ III , Arkiv för Mat. (1969), 509 - 512.
- A6. G. Aronsson, On Certain Singular Solutions of the Partial Differential Equation $u_x^2 u_{xx} + 2 u_x u_y u_{xy} + u_y^2 u_{yy} = 0$, Manuscripta Math. 47 (1984), 133 - 151.
- A7. G. Aronsson, Construction of Singular Solutions to the p -Harmonic Equation and its Limit

- FL. I. Fonseca, G. Leoni, *Modern methods in the Calculus of Variations: L^p spaces*, Springer Monographs in Mathematics, 2007.
- HJ. R.A. Horn, Ch.R. Johnson, *Matrix Analysis*, Cambridge University Press, 2012.
- K1. N. Katzourakis, L^1 -Variational Problems for Maps and the Aronsson PDE system, *J. Differential Equations* 253 (2012), 2123 - 2139.
- K2. N. Katzourakis, L^1 -Minimal Submanifolds, *Proceedings of the AMS* 142 (2014), 2797 - 2811.
- K3. N. Katzourakis, On the Structure of L^1 -Harmonic Maps, *Communications in PDE* 39 (2014), 2091 - 2124.
- K4. N. Katzourakis, Explicit 2D L^1 -Harmonic Maps whose Interfaces have Junctions and Corners, *Comptes Rendus Acad. Sci. Paris, Ser. I* 351 (2013), 677 - 680.
- K5. N. Katzourakis, Optimal L^1 -Quasiconformal Immersions, *ESAIM Control, Opt. and Calc. Var.*, to appear (2015) DOI: <http://dx.doi.org/10.1051/cocv/2014038>.
- K6. N. Katzourakis, Nonuniqueness in Vector-valued Calculus of Variations in L^1 and some Linear Elliptic Systems, *Comm. on Pure and Appl. Anal.* 14 (2015), 313 - 327.
- K7. N. Katzourakis, An Introduction to viscosity Solutions for Fully Nonlinear PDE with Applications to Calculus of Variations in L^1 , *Springer Briefs in Mathematics*, 2015, DOI 10.1007/978-3-319-12829-0.
- K8. N. Katzourakis, Generalised solutions for fully nonlinear PDE systems and existence-uniqueness theorems, *ArXiv preprint*, <http://arxiv.org/pdf/1501.06164.pdf>.
- K9. N. Katzourakis, Absolutely minimising generalised solutions to the equations of vectorial Calculus of Variations in L^1 , *ArXiv preprint*, <http://arxiv.org/pdf/1502.01179.pdf>.
- K10. N. Katzourakis, A new characterisation of L^1 -Harmonic and p -Harmonic maps via a new variations in L^1 , *ArXiv preprint*, <http://arxiv.org/pdf/1509.01811.pdf>.
- K11. N. Katzourakis, Minimization of D -solutions to fully nonlinear PDE systems, *ArXiv preprint*, <http://arxiv.org/pdf/1508.05519.pdf>.
- K12. N. Katzourakis, Equivalence between weak and D -solutions for symmetric hyperbolic first order PDE systems, *ArXiv preprint*, <http://arxiv.org/pdf/1507.03042.pdf>.
- K13. N. Katzourakis, Solutions of vectorial Hamilton-Jacobi equations are rank-one Absolute Minimisers in L^1 , *ArXiv preprint*, <http://arxiv.org/abs/1604.00802>.
- KP. N. Katzourakis, T. Pryer, On the numerical approximation of L^1 -Harmonic mappings, *ArXiv preprint*, <http://arxiv.org/pdf/1511.01308.pdf>.
- P. P. Pedregal, *Parametrized Measures and Variational Principles*, Birkhäuser, 1997.
- Pi. G. Pisante, Sufficient conditions for the existence of viscosity solutions for nonconvex Hamiltonians, *SIAM J. Math. Anal.*, 36(1):186203, 2004.
- V. M. Valadier, Young measures, in "Methods of nonconvex analysis", *Lecture Notes in Mathematics* 1446, 152-188 (1990).

Normandie Univ, UNIHAVRE, LMAH, 76600 Le Havre, France
E-mail address : gisella.croce@univ-lehavre.fr

Department of Mathematics and Statistics, University of Reading, Whiteknights,
PO Box 220, Reading RG6 6AX, UK
E-mail address : n.katzourakis@reading.ac.uk

Dipartimento di Matematica e Fisica, Seconda Università degli Studi di Napoli -
S.U.N., Viale Lincoln, 5, c.a.p. 81100 Caserta, Italy
E-mail address : giovanni.pisante@unina2.it